

# Math 210A Lecture 6 Notes

Daniel Raban

October 10, 2018

## 1 Inverse Limits, Direct Limits, and Adjoint Functors

### 1.1 Inverse and direct limits

**Example 1.1.** Consider the colimit of this diagram in  $\text{Ab}$ :

$$\mathbb{Z}/p\mathbb{Z} \xrightarrow{p} \dots \xrightarrow{p} \mathbb{Z}/p^n\mathbb{Z} \xrightarrow{p} \mathbb{Z}/p^{n+1}\mathbb{Z} \xrightarrow{p} \dots$$

Then  $\varinjlim_n \mathbb{Z}/p^n\mathbb{Z} \cong \mathbb{Q}_p/\mathbb{Z}_p \subseteq \mathbb{Q}/\mathbb{Z}$ , where  $\mathbb{Q}_p$  is the free field of  $\mathbb{Z}_p$ . We can also show that  $\mathbb{Q}_p/\mathbb{Z}_p = \{a \in \mathbb{Q}/\mathbb{Z} : p^n a = 0 \text{ for some } n \geq 0\}$ .

**Definition 1.1.** A **directed set**  $I$  is a set with a partial ordering such that for all  $i, j \in I$ , there is a  $k \in I$  such that  $i \leq k, j \leq k$ .


**Definition 1.2.** A **directed category** is a category where the objects are elements of a directed set  $I$ , and there are morphisms  $i \rightarrow j$  iff  $i \leq j$ . A **codirected category**  $\mathcal{I}$  is a category where  $\mathcal{C}^{op}$  is directed.

**Definition 1.3.** Suppose  $\mathcal{I}$  is codirected with  $\text{Obj}(\mathcal{I}) = I$  and  $F : \mathcal{I} \rightarrow \mathcal{C}$ . A limit of  $F$  is called the **inverse limit** of the  $F(i)$  for all  $i \in I$ . We write  $\lim F = \varprojlim_{i \in I} F(i)$ .

$$\begin{array}{ccc} & & F(i) \\ & \nearrow & \nearrow \\ \varprojlim F(i) & \longrightarrow & F(k) \\ & \searrow & \searrow \\ & & F(j) \end{array}$$

If  $\mathcal{I}$  is directed with  $\text{Obj}(\mathcal{I}) = I$  and  $F : \mathcal{I} \rightarrow \mathcal{C}$ . A colimit of  $F$  is called the **direct limit**

of the  $F(i)$  for all  $i \in I$ . We write  $\text{colim } F = \varinjlim_{i \in I} \text{colim } F$ .



**Definition 1.4.** A small category  $\mathcal{I}$  is **filtered** if

1. for all  $i, j \in I$ , there exists  $k \in I$  such that there exist morphisms  $i \rightarrow k, j \rightarrow k$ ,
2. for all  $\kappa, \kappa' : i \rightarrow j$  in  $I$  there exists a morphism  $\lambda : j \rightarrow k$  such that  $\lambda \circ \kappa = \lambda \circ \kappa'$

A category is **cofiltered** if the opposite category is filtered.

Cofiltered limits and filtered limits generalize inverse and direct limits, respectively.

**Example 1.2.** Let  $I$  be cofiltered with an initial object  $c$ . Then if  $F : I \rightarrow \mathcal{C}$ ,  $\lim F = F(c)$ .

## 1.2 Adjoint functors

**Definition 1.5.** A functor  $F : \mathcal{C} \rightarrow \mathcal{D}$  is **left adjoint** to a functor  $G : \mathcal{D} \rightarrow \mathcal{C}$  if for each  $C \in \mathcal{C}, D \in \mathcal{D}$ , there exist bijections  $\eta_{C,D} : \text{Hom}_{\mathcal{D}}(F(C), D) \rightarrow \text{Hom}_{\mathcal{C}}(C, G(D))$  such that  $\eta$  is a natural transformation between functors  $\mathcal{C}^{op} \times \mathcal{D} \rightarrow \text{Sets}$ . That is,

$$\begin{array}{ccc} \text{Hom}_{\mathcal{D}}(F(C), D) & \xrightarrow{\eta_{C,D}} & \text{Hom}_{\mathcal{C}}(C, G(D)) \\ \downarrow h \mapsto g \circ h \circ F(f) & & \downarrow h \mapsto G(g) \circ h \circ f \\ \text{Hom}_{\mathcal{D}}(F(C'), D') & \xrightarrow{\eta_{C',D'}} & \text{Hom}_{\mathcal{C}}(C', G(D')) \end{array}$$

$G$  is **right adjoint** to  $F$  if  $F$  is left adjoint to  $G$ .

**Remark 1.1.** If  $F : \mathcal{C} \rightarrow \mathcal{D}$  and  $G : \mathcal{D} \rightarrow \mathcal{C}$  are quasi-inverses and  $\eta : \text{id}_{\mathcal{C}} \rightarrow G \circ F$  is a natural isomorphism, then we can define  $\phi_{C,D} : \text{Hom}_{\mathcal{D}}(F(C), D) \rightarrow \text{Hom}_{\mathcal{C}}(C, G(D))$  given by  $h \mapsto G(h) \circ \eta_C$ . Check that  $\phi_{C,D}$  is a bijection. So  $F$  is left-adjoint to  $G$ . Similarly,  $G$  is left-adjoint to  $F$ .

**Proposition 1.1.** Suppose  $S$  is a set, and consider  $h_S : \text{Set} \rightarrow \text{Set}$  given by  $h_S(T) = \text{Maps}(S, T)$  and  $h_S(f : T \rightarrow T') = g \mapsto f \circ g$ . Then  $h_S$  is right adjoint to  $t_S : \text{Set} \rightarrow \text{Set}$  given by  $t_S(T) = T \times S$  and  $t_S(f) = (f, \text{id}_S) : T \times S \rightarrow T' \times S$ .

*Proof.* We need to find a bijection  $\tau_{T,U} : \text{Maps}(T \times S, U) \rightarrow \text{Maps}(T, \text{Maps}(S, U))$ . We can send  $f \mapsto (t \mapsto (s \mapsto f(s, t)))$ . To show that this is a bijection, we can go backward by sending  $\varphi \mapsto ((t, s) \mapsto \varphi(t)(s))$ . Check that these maps are inverses of each other and that this is a natural transformation.  $\square$

**Proposition 1.2.** *Suppose all limits  $F : I \rightarrow \mathcal{C}$  exist. Then the functor  $\lim : \text{Fun}(I, \mathcal{C}) \rightarrow \mathcal{C}$  given by  $F \mapsto \lim F$  and  $(\eta : F \rightarrow F') \mapsto (\lim F \mapsto \lim F')$  has a left adjoint  $\Delta : \mathcal{C} \rightarrow \text{Fun}(I, \mathcal{C})$  such that  $\Delta(A) = c_A$  is the constant functor  $I \rightarrow \mathcal{C}$  with value  $A$ .*

*Proof.* We want a bijection  $\eta : \text{Hom}_{\text{Fun}(I, \mathcal{C})}(c_A, F) \rightarrow \text{Hom}_{\mathcal{C}}(A, \lim F)$ . Let  $\eta : c_A \rightarrow F$  be  $\eta_i : \underbrace{c_A(i)}_{=A} \rightarrow F(i)$  such that

$$\begin{array}{ccc} A & \xrightarrow{\eta_i} & F(i) \\ \text{id}_A = c_A(f) \downarrow & & \downarrow F(f) \\ A & \xrightarrow{\eta_j} & F(j) \end{array} \qquad \begin{array}{ccc} A & \xrightarrow{\eta_i} & F(i) \\ & \searrow \eta_j & \downarrow F(f) \\ & & F(j) \end{array}$$

for all  $f : i \rightarrow j$ . So  $\eta_j = F(f) \circ \eta_i$  for all  $f : i \rightarrow j$ . There exists a unique morphism  $g : A \rightarrow \lim F$  such that

$$\begin{array}{ccc} & A & \\ & \swarrow \eta_j & \searrow \eta_i \\ & \lim F & \\ & \swarrow p_j & \searrow p_i \\ F(j) & \xleftarrow{F(f)} & F(i) \end{array}$$

Send  $\eta$  to  $g$ . Conversely if we have  $g : A \rightarrow \lim F$ ,  $\eta_i = p_i \circ g$  is a morphism from  $A \rightarrow F(i)$ . So we get  $\eta \in \text{Hom}_{\text{Fun}(I, \mathcal{C})}(c_A, F)$ .  $\square$

**Definition 1.6.** A contravariant functor  $F : \mathcal{C} \rightarrow \text{Set}$  is **representable** if there exists an object  $B \in \mathcal{C}$  and a natural isomorphism  $h^B \rightarrow F$ , where  $h^B = \text{Hom}_{\mathcal{C}}(\cdot, B)$ . We say that  $B$  **represents**  $F$ .

**Example 1.3.** The functor  $P : \text{Set} \rightarrow \text{Set}$  given by  $S \mapsto \mathcal{P}(S)$  and  $(f : S \rightarrow T) \mapsto (V \mapsto f^{-1}(V))$  is representable by  $\{0, 1\}$ .