

# Math 255A Lecture 9 Notes

Daniel Raban

October 17, 2018

## 1 Applications of Baire's Theorem I: The Open Mapping Theorem

### 1.1 The open mapping theorem

Banach used Baire's theorem to prove a number of striking results in functional analysis. Recall Baire's theorem.

**Theorem 1.1** (Baire category). *Let  $E$  be a complete metric space, and let  $(F_n)_{n \in \mathbb{N}}$  be closed in  $E$  containing no interior points. Then the union  $\bigcup_{n=1}^{\infty} F_n$  has no interior points either. Moreover,  $E \neq \bigcup_{n=1}^{\infty} F_n$ .*

**Definition 1.1.** We say that  $A \subseteq E$  is **of the first category** (or **meager**) if there exists a sequence  $F_n$  of closed sets without interior points such that  $A \subseteq \bigcup_{n=1}^{\infty} F_n$ .

**Theorem 1.2** (Banach, open mapping theorem). *Let  $F_1, F_2$  be Fréchet spaces, and let  $T : F_1 \rightarrow F_2$  be linear continuous. Then either  $\text{im}(T) \subseteq F_2$  is of the first category, or else  $\text{im}(T) = F_2$  and the mapping  $T$  is open.*

*Proof.* Let  $U$  be an open neighborhood of 0 in  $F_1$ . We claim that  $\overline{T(U)}$  contains a neighborhood of 0 in  $F_2$ , provided  $\text{im}(T)$  is not of the first category. Let  $V$  be a balanced neighborhood of 0 in  $F_1$  such that  $V + V \subseteq U$ . Then  $V$  is absorbing (for  $x \in F_1$ ,  $\lambda x \in V$  for sufficiently small  $|\lambda|$ ). So  $F_1 = \bigcup_{n=1}^{\infty} nV$  means that  $\text{im}(T) = \bigcup_{n=1}^{\infty} T(nV) \subseteq \bigcup_{n=1}^{\infty} \overline{T(nV)}$ . Since  $\text{im}(T)$  is not of the first category, for some  $n$ ,  $\overline{T(nV)} = n\overline{T(V)}$  has an interior point. Then  $\overline{T(V)}$  has an interior point. So there exists  $y \in F_2$  and a neighborhood  $W$  of 0 in  $F_2$  such that  $\{y\} + W \subseteq \overline{T(V)}$ . Then  $y \in \overline{T(V)}$ .  $V = -V$  since  $V$  is balanced, so  $-y \in \overline{T(V)}$ . So  $W \subseteq \overline{T(V)} + \{-y\} \subseteq \overline{\overline{T(V)} - T(V)} = \overline{T(V) - T(V)}$ . We get  $W \subseteq \overline{T(V+V)} \subseteq \overline{T(U)}$ , as claimed.

Let  $d_{F_1}$  be a translation invariant metric on  $F_1$  generating the topology on  $F_1$ , and define  $d_{F_2}$  similarly. Thus, for any  $r > 0$ , there exists  $\rho > 0$  such that  $B_{F_2}(0, \rho) \subseteq T(B_{F_1}(0, r))$ . The metrics  $d_{F_1}, d_{F_2}$  are translation invariant, so for any  $r > 0$ , there exists a  $\rho > 0$  such that for any  $x \in F_1$ ,  $B_{F_2}(Tx, \rho) \subseteq \overline{T(B_{F_1}(x, r))}$ . Let  $r > 0$  be arbitrary and let  $r_n = r/2^n$

for  $n \in N$ . We get the corresponding  $\rho_n$  sequence such that  $B_{F_2}(Tx, \rho_n) \subseteq \overline{T(B_{F_1}(x, r_n))}$  for all  $x \in F_1$ . We can arrange so that  $\rho_n \downarrow 0$ .

Let  $y \in B_{F_2}(Tx, \rho_0)$ . We shall show that there is an  $x' \in F_1$  such that  $d_{F_1}(x, x') \leq 2r$  and  $y = Tx'$ . Let  $x_1 \in \overline{B_{F_1}(x, r_0)}$  be such that  $d_{F_2}(y, Tx_1) < \rho_1 \iff y \in B_{F_2}(Tx_1, \rho_1) \subseteq \overline{T(B_{F_1}(x_1, r_1))}$ . Let  $x_2 \in \overline{B_{F_1}(x_1, r_1)}$  be such that  $d_{F_2}(y, Tx_2) < \rho_2$ . Then  $y \in B_{F_2}(Tx_2, \rho_2) \subseteq \overline{T(B_{F_1}(x_2, r_2))}$ . Continuing in this fashion, we get a sequence  $(x_n)$  in  $F_1$  such that  $x_{n+1} \in \overline{T(B_{F_1}(x_n, r_n))}$ . Then  $(x_n)$  is a Cauchy sequence in  $F_1$ , and  $d_{F_2}(y, Tx_n) < \rho_n \rightarrow 0$ . We get  $x_n \rightarrow x' \in F_1$ , where  $d_{F_1}(x, x') \leq 2r$ , and, since  $T$  is continuous,  $Tx_n \rightarrow Tx'$ . So  $y = Tx'$ .

So we get that for all  $r > 0$ , there exists  $\rho > 0$  such that  $B_{F_2}(Tx, \rho) \subseteq T(B_{F_1}(x, 2r))$ . Hence,  $\text{im}(T) = F_2$ , and  $T$  is open.  $\square$

**Corollary 1.1.** *Let  $T : F_1 \rightarrow F_2$  be an injective, linear, continuous map between Fréchet spaces. Then either the range of  $T$  is of the first category, or  $\text{im}(T) = F_2$ , and  $T$  is a homeomorphism.*

## 1.2 Application of the open mapping theorem to partial differential equations

Let  $P(D) = \sum_{|\alpha| \leq m} a_\alpha D^\alpha$ , where  $D^\alpha = D_{x_1}^{\alpha_1} \cdots D_{x_n}^{\alpha_n}$  and  $D_{x_j} = (1/i)\partial_{x_j}$  be a partial differentiation operator (on  $\mathbb{R}^n$ ) with constant coefficients  $a_\alpha \in \mathbb{C}$ . Assume that for some open set  $\Omega \subseteq \mathbb{R}^n$ , every solution  $u \in C^m(\Omega)$  of  $Pu = 0$  is in fact in  $C^{m+1}(\Omega)$  (e.g.  $P = \Delta$ , the Laplacian). Then we have  $\text{Im}(\zeta) \rightarrow \infty$  if  $\zeta \rightarrow \infty$  on the surface in  $\mathbb{C}^n$  given by  $0 = P(\zeta) = \sum_{|\alpha| \leq m} a_\alpha \zeta^\alpha$ . We will do this in detail next time.